

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 December 2000 (21.12.2000)

PCT

(10) International Publication Number
WO 00/76955 A1

- (51) International Patent Classification⁷: C07C 215/52, A61K 31/135, A61P 25/24
- (21) International Application Number: PCT/US00/16388
- (22) International Filing Date: 14 June 2000 (14.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/333,594 15 June 1999 (15.06.1999) US
- (71) Applicant: AMERICAN HOME PRODUCTS CORPORATION [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US).
- (72) Inventors: YARDLEY, John, Patrick; 154 Hughes Road, King of Prussia, PA 19406 (US). ASSELIN, Andre, Alfred; 64 Ackerman Drive, Mahwah, NJ 07430 (US).
- (74) Agents: ECK, Steven, R.; American Home Products Corporation, Patent Law Dept. - 2B, One Campus Drive, Parsippany, NJ 07054 et al. (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 00/76955 A1

(54) Title: ENANTIOMERS OF O-DESMETHYL VENLAFAXINE

(57) Abstract: This invention provides pharmaceutically active enantiomers of the venlafaxine metabolite O-Desmethyl venlafaxine, R(-)-4-[2-(Dimethylamino)-1-(1-hydroxycyclo-hexyl)ethyl]phenol or R(-)-1-[2-(dimethylamino)-1-(4-hydroxyphenyl)-ethyl]cyclo-hexanol, and S(+)-1-[2-(Dimethylamino)-1-(4-hydroxyphenyl)ethyl]-cyclohexanol or S(+)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol, or one or more pharmaceutically acceptable salts or salt hydrates thereof, as well as pharmaceutical compositions utilizing these enantiomers and methods of using the enantiomers to treat, inhibit or control central nervous system disorders.

- 1 -

ENANTIOMERS OF O-DESMETHYL VENLAFAXINE

This invention provides enantiomers of O-desmethyl venlafaxine, (R/S) 4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol, as well as pharmaceutical compositions and uses thereof.

Background of the Invention

Various patents and literature references describe the biological activities of venlafaxine, and its salts and analogs. Venlafaxine hydrochloride tablets are marketed by Wyeth-Ayerst Laboratories under the Effexor® trademark.

The absolute configuration of the (+) enantiomer of venlafaxine was established as S by a single crystal X-ray analysis of the hydrobromide salt and the anomalous dispersion technique (Yardley et al., J. Med. Chem., 1990, 33, 2899).

(R/S)-1-[2-(dimethylamino)-1-(4-methoxyphenyl)ethyl]cyclohexanol and its metabolites 1-[2-(dimethylamino)-1-(4-hydroxyphenyl)ethyl]cyclohexanol and 1-[1-(4-methoxyphenyl)-2-(methylamino)ethyl]cyclohexanol are disclosed and claimed in U.S. Patent No. 4,535,186 (Husbands et al.). U.S. Patent No. 5,530,013 (Husbands et al.) claims the use of venlafaxine in the inducement of cognition enhancement. U.S. Patent No. 5,506,270 (Upton et al.) claims venlafaxine's use in methods of treating hypothalamic amenorrhea in non-depressed women.


U.S. Patents Nos. 5,788,986 (Dodman) and 5,554,383 (Dodman) teaches and claims the use of serotonin reuptake inhibitors in modifying the behavior of dogs.

Summary of the Invention

This invention provides pharmaceutically active enantiomers of the venlafaxine metabolite O-Desmethyl venlafaxine, R(-)-4-[2-(Dimethylamino)-1-(1-hydroxycyclo-hexyl)ethyl]phenol and S(+)-4-[2-(Dimethylamino)-1-(1-hydroxy-

- 2 -

cyclo-hexyl)ethyl]-phenol, or a pharmaceutically acceptable salt or salt hydrate thereof, having the structures:



Particularly, this invention provides both the R(-) enantiomer and S(+)
 5 enantiomer substantially free of each other. Under a different system of nomenclature S(+)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)-ethyl]phenol may also be named S(+)-1-[2-(Dimethylamino)-1-(4-hydroxyphenyl)-ethyl]cyclohexanol. Similarly, R(-)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)-ethyl]phenol may also be referred to as R(-)-1-[2-(dimethylamino)-1-(4-hydroxyphenyl)-ethyl]cyclohexanol.
 10 As used herein, the designations (+) and (-) refer to the sign of rotation of the relevant free base.

These enantiomers and their pharmaceutically useful salts and hydrates are useful for the biological and pharmacological activities for which venlafaxine and its salts are known in the art. These enantiomers may be used in treating or inhibiting
 15 central nervous system disorders, including depression, panic disorder, post-traumatic stress disorder, late luteal phase dysphoric disorder (also known as pre-menstrual syndrome), attention deficit disorder, with and without hyperactivity, generalized anxiety disorder, bulimia nervosa, Gilles de la Tourette Syndrome, Shy Drager Syndrome, vasomotor flushing, drug and alcohol addiction, sexual dsifunction
 20 (including premature ejaculation), borderline personality disorder, chronic fatigue syndrome, fibromyalgia, urinary incontinence and others. These compounds are also useful in the inducement of cognition enhancement and in regimens for cessation of smoking or other tobacco uses.

Racemic 1-[2-(dimethylamino)-1-(4-hydroxyphenyl)ethyl]cyclohexanol can
 25 be produced as described in Example 26 of U.S. Patent No. 4,535,186 (Husbands et

- 3 -


al.), which is incorporated herein by reference. It will be understood that the enantiomers may be separated from each other by standard resolution techniques known in the art. For example a racemic mixture may be converted to a mixture of optically active diastereoisomers by reaction with a single enantiomer of a 'resolving agent' (for example by diastereomeric salt formation or formation of a covalent bond).
 5 The resulting mixture of optically active diastereoisomers may be separated by standard techniques (e.g. crystallisation or chromatography) and individual optically active diastereoisomers then treated to remove the 'resolving agent' thereby releasing the single enantiomer of the compound of the invention. Chiral chromatography
 10 (using a chiral support, eluent or ion pairing agent) may also be used to separate enantiomeric mixtures directly

Alternatively, these R and S enantiomers may be obtained by subjecting R-4-[2-(dimethylamino)-1-(1,-R₁O cyclohexyl)ethyl]phenol or a salt thereof or (S)-4-[2-(dimethylamino)-1-(1,-R₁O cyclohexyl)ethyl]phenol or a salt thereof, wherein R₁ is a
 15 removable protecting group, to a reaction to remove the protecting group. The group R₁O may be, for example, benzyloxy or methoxy. The benzyl group may be removed by reduction, particularly by hydrogenation. In the case where R₁O is methoxy the R and S enantiomers may be obtained by O-demethylation of the
 20 separated enantiomers of venlafaxine using, for example, either boron tribromide or ethane thiol anion.

Example No. 1

1-[2-(Dimethylamino-1-(4-hydroxyphenyl)ethyl)cyclohexanol fumarate hydrate

25




- 4 -

1-[2-(Dimethylamino-1-(4-methoxyphenyl)ethyl)cyclohexanol -HCl (200 g = 0.6372 mol) was dissolved in H₂O (500 mL). CH₂Cl₂ (350 mL) was added thereto and this mixture cooled to 10°C. At this temperature 2.5 N NaOH (280 mL= 0.7 mol) was added slowly over 1 hr. CH₂Cl₂ was separated and the aqueous layer extracted with CH₂Cl₂ (200 mL). Combined CH₂Cl₂ extracts were dried (MgSO₄) filtered, and evaporated in vacuo to yield 1-[2-(Dimethylamino-1-(4-methoxyphenyl)-ethyl)cyclohexanol free base (167.5 g = 94.5%) as white solid, mp 77-79°C.

10 To a stirred solution of 1-[2-(Dimethylamino-1-(4-methoxyphenyl)ethyl)-cyclohexanol -free base (13.87 g = 50 mmols) in CH₂Cl₂ (300 mL) cooled to -40°C, under N₂ was added slowly BBr₃ (10 mL = 105.5 mmols) over a period of 15 minutes. The reaction mixture warmed to 0°C where it was stirred for 3 hrs. During this time a gummy precipitate formed. Still at 0°C, 2.5N NaOH (200 mL) was added slowly over 1 hr, then allowed to warm to room temperature and stirred for 3 hrs. CH₂Cl₂ was removed by evaporation under reduced pressure leaving an aqueous layer having a pH = 13-14. Aqueous layer was extracted with EtOAc (3 x 100 mL) and its pH dropped to 9. Combined EtOAc extracts were dried (MgSO₄), filtered, evaporated in vacuo to afford crude phenol (9.3 g = 71%) as a white solid, mp 208-213°C (TLC) together with some dehydrated product. This crude product was used in the next step without further purification.

25 Crude phenol (9.3 g = 35.31 mmoles) and fumaric acid (4.91 g = 42.37 mmoles) were dissolved in a mixture of methanol/acetone (1:3) (195 mL), stirred at room temperature for 15 minutes. H₂O (0.8 mL = 44.44 mmoles) was added to the clear light yellow solution. The whole was stirred at room temperature for 3 hours. The resulting white precipitate was filtered, washed with acetone (30 mL) dried in air to yield 8.0 g = 57% white solid, mp: 145-150°C.

- 5 -

Example No. 2**R(-)-1-[2-(Dimethylamino-1-(4-methoxyphenyl)ethyl)cyclohexanol**

To a solution of yield 1-[2-(Dimethylamino-1-(4-methoxyphenyl)ethyl)-cyclohexanol free base (100 g = 0.36 mol) in EtOAc (750 mL) at room temp was added at once a solution of (+)-Di-para Toluoyl-D-tartaric acid-monohydrate (DT(-)T; 40 g = 0.0991 mol). The whole was stirred at room temp for 1 hr. The resulting precipitate was filtered off, washed with EtOAc (3 x 100 mL), dried overnight at 35°C in a vacuum oven to provide crude R(-)-1-[2-(dimethylamino)-1-(4-methoxyphenyl)-ethyl]cyclohexanol DT(-)T salt (83 g = 92.8%) as a white solid.

15 **Recrystallization of R(-)-1-[2-(Dimethylamino-1-(4-methoxyphenyl)-ethyl]cyclohexanol, DT(-)T Salt**

Crude DT(-)T salt (83g) was dissolved in EtOAc (700 mL). The mixture was heated up to reflux. Methanol (75 mL) was added thereto to obtain a clear solution. The mixture was concentrated at atmospheric pressure to a volume of 400 mL (some solid started to precipitate). The mixture was cooled at 25°C for 1 hr, then at 0°C for another 2 hrs and filtered off to provide (-)-1-[2-(Dimethylamino-1-(4-methoxyphenyl)ethyl]cyclohexanol DT(-)T salt (63 g=77%). NOTE: Optical rotation of this salt in ethanol was + 47.0°.


20

- 6 -

Isolation of R(-)-1-[2-(Dimethylamino-1-(4-methoxyphenyl)-ethyl]cyclohexanol Base

R(-)-1-[2-(Dimethylamino-1-(4-methoxyphenyl)ethyl]cyclohexanol DT(-)T
 5 salt was slurried in a mixture of H₂O/CH₂Cl₂ (400 mL/400mL). The pH of this mixture was adjusted to 13 by adding 25% NaOH solution (120 mL). The layers were separated, aqueous layer was extracted with CH₂Cl₂ (1 x 200 mL). Combined CH₂Cl₂ layers were washed with H₂O (2 x 200 mL) saturated NaCl solution (1 x 200 mL), dried (MgSO₄) and evaporated at atmospheric pressure to a volume of 100 mL.
 10 Hexane (300 mL) was added thereto and solution became hazy. After charcoal treatment (1 teaspoon), the filtrate was concentrated at atmospheric pressure to a volume of 250 mL and allowed to cool. The resulting white precipitate was collected by filtration, washed well with hexane (2 x 100 mL), dried in a vacuum oven overnight to provide R(-)-1-[2-(Dimethylamino-1-(4-methoxyphenyl)ethyl]cyclohexanol - base (28.5 g). Recrystallization from CH₂Cl₂/hexane (50 mL/200 mL) gave analytically pure R(-)-1-[2-(Dimethylamino-1-(4-methoxyphenyl)ethyl]cyclohexanol base (23.5 g = 23.5%) Rotation = -31.08° (in ethanol). Anal. Calcd: C, 73.60; H, 9.81; N, 5.05 Found: C, 73.75, H, 9.73; N, 4.83.

20

Example No. 3**R(-)-1-[2-(Dimethylamino-1-(4-hydroxyphenyl)ethyl]cyclohexanol fumarate hydrate salt**


25

To a cooled (-40°C) stirred solution of R(-)-1-[2-(dimethylamino)-1-(4-methoxyphenyl) ethyl]cyclohexanol (13.87 g = 50 mmol) in CH₂ Cl₂ (500 mL) under nitrogen was added slowly BBr₃ (10 mL = 105.5 mmols) over a period of 15 min. The reaction mixture warmed to 0°C where it was stirred for 3 h. During this time a

- 7 -

gummy precipitate formed. Still at 0°C, 2.5 N NaOH solution (200 mL) was added slowly over 1 hr. The reaction mixture was allowed to warm to room temp and stirred overnight. Methylenechloride was removed, leaving an aqueous layer having a pH - 13-14. Aqueous layer was extracted with EtOAc (3 x 100 mL) and its pH
5 dropped in 9. Combined EtOAc extracts were washed with brine, dried (MgSO₄) and evaporated in vacuo to give crude phenol (6.5 g - 49.4%) as white solid. The crude phenol (6.5 g = 24.68 mmols) and fumaric acid (1.2 eq; 3.3 g = 29.61 mmols) were dissolved in a mixture of methanol/acetone (1:3) (135 mL) and stirred at room temp for 15 min. After this time H₂O (0.6 mL) was added to the clear light yellow colored
10 solution. Precipitation was seen immediately. The suspension was stirred at room temp for 3 h. The resulting white precipitate was collected by filtration, washed well with acetone (1 x 35 mL) and dried to provide title compound (6.6 g = 67.3%, mp 150-155°C. This was recrystallized from MeOH/Acetone/H₂O (40 mL: 126 mL: 0.3 mL) to give 3.7 g (37.7%) of analytically pure R(-)-4-[2-(Dimethylamino)-1-(1-
15 hydroxycyclo-hexyl)ethyl]phenol fumarate hydrate salt. Rotation: + 6.60° (in methanol) for the fumarate hydrate, -15.58° (in methanol) for the free base. Anal. Calcd: C, 60.43; H, 7.86; N, 3.52. Found: C, 60.16; H, 7.64; N, 3.36.

- 8 -

Example No. 4**R(-)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol fumarate
hydrate salt**

5

Under gentle N_2 stream 60% NaH (35 mmols=1.4 g) was washed with hexane, collected by filtration and transferred into a 250 mL 3 neck flask. DMF (20 mL) was added into the flask to cover the sodium hydride and the suspension cooled to 10°C .

- 10 Under stirring a solution of ethane thiol (32.40 mmols=2.075 g=2.47 mL) in DMF (5 mL) was added dropwise over 40 min. During the addition of the ethanethiol solution, foaming was noted in the flask. Stirring was continued at 20°C for 1 1/2 h and the starting material R(-)-1-[2-(dimethylamino)-1-(4-methoxyphenyl)ethyl]cyclohexanol (12.5 mmols = 3.46 g) was added as a solid over 5 min. The reaction
- 15 mixture was heated up slowly to 150°C over 30 min. and stirred at this temperature for another hour. After this time the yellow brownish colored reaction mixture was rapidly cooled to 25°C and quenched by adding it into a flask containing H_2O (90mL).

- 9 -

The mixture was charcoaled and filtered through celite. The mixture was washed with H₂O (1 x 25 mL) and 1N NaOH (1 x 50 mL). At this point the pH of the clear yellow colored solution was 13. This was extracted with toluene (1 x 60 mL), followed by hexane (1 x 60 mL). Under stirring at room temp it was neutralized to pH=9 with conc. HCl (3 mL). The resulting suspension was cooled at 5°C, stirred for 1h and the white solid was collected by filtration, dried in air overnight to give 2.6 g=79% of the phenol (mp:232-235°C). This compound (9.491 mmols=2.5 g) and fumaric acid (11.39 mmols=1.32 g) were dissolved in hot methanol/acetone (1:3) mixture (54 mL) and filtered leaving a clear light yellow colored solution. Under stirring at room temp H₂O (0.227 mL) was added to the solution. The solution became cloudy immediately. Stirring was resumed for 2h and the resulting white solid was collected by filtration, washed with acetone (2 x 50 mL), dried at 35°C in a vacuum oven overnight to give 3 g=60% of analytically pure compound.

15

Example No. 5**S(+)-1-[2-(Dimethylamino)-1-(4-hydroxyphenyl)ethyl]cyclohexanol fumarate hydrate salt**

a) S(+)-1-[2-(Dimethylamino)-1-(4-methoxyphenyl)ethyl]cyclohexanol

20


To the mother liquor from the resolution after separation of R(-)-1-[2-(dimethylamino)-1-(4-methoxyphenyl)ethyl]cyclohexanol DT(-)T salt (see Example No. 2) was added H₂O (400 mL). The mixture has a pH=7. The pH was adjusted to 12 by adding 25% NaOH solution (150 mL). EtOAc layer was separated, washed with saturated NaCl solution (2 x 100 ml) dried (MgSO₄) and concentrated in vacuo to a volume of 100 mL. Hexane (400 mL) was added thereto and the whole was stirred at room temp for 1 h, then at 0°C for another 2 h. The white precipitate was collected by filtration to give 37.5 g. This was dissolved in hot CH₂Cl₂ (110 mL). Charcoal (2 g) was added to the hot solution and stirred for 5 minutes. After filtration through solka floc, hexane (380 mL) was added to the filtrate. The mixture was concentrated at atmospheric pressure to a volume of 250 mL and allowed to stay at room temp overnight. The resulting white precipitate was collected by filtration to provide 29.7 g. Recrystallization from CH₂Cl₂/Hexane (72/249 mL) gave analytically

25
30

- 10 -

pure S(+)-1-[2-(dimethylamino)-1-(4-methoxyphenyl)ethyl]-cyclohexanol, 25.4 g = 25.4 % yield. Rotation: +28.82° (in ethanol.). Anal. Calcd.: C, 73.60; H, 9.81; N, 5.05. Found: C, 73.70; H, 10.10; N, 4.85.

- 5 b) S(+)-1-[2-(Dimethylamino)-1-(4-hydroxyphenyl)ethyl]cyclohexanol fumarate hydrate salt



- 10 To a stirred solution of S(+)-1-[2-(dimethylamino)-1-(4-methoxyphenyl)-ethyl]-cyclohexanol (13.87 g = 50 mmols) in CH₂Cl₂ (500mL), cooled to -40°C under nitrogen was added slowly BBr₃ (10mL = 105.5 mmols) over a period of 15 min. The reaction mixture warmed to 0°C where it was stirred for 3 h. During this time a gummy precipitate formed. Still at 0°C, 2.5 N NaOH (200mL) was added slowly
- 15 over 1 h. Then the mixture was allowed to warm to room temp and stirred overnight. CH₂Cl₂ was removed under vacuo leaving an aqueous layer having a pH=13-14. Aqueous layer was extracted with EtOAc (3 × 100 mL) and its pH dropped to 9. Combined EtOAc extracts were washed with saturated NaCl solution, dried (MgSO₄), filtered and evaporated in vacuo to afford crude phenol (3.9 g = 29.6%) as a white
- 20 solid. Crude phenol (3.9 g = 14.8 mmols) and fumaric acid (2.06 g = 17.77 mmols) were dissolved in a mixture of methanol/acetone (1:3) (81 mL) and stirred at room temp for 15 min. H₂O (0.325 mL = 18 mmol) was added to the clear light yellow solution. Precipitation was noted immediately. The whole was stirred at room temp for 3 h. The resulting white precipitate was collected by filtration, washed with
- 25 acetone (1 × 35 mL) and dried to give 2.4 g (40.8%) of product. Recrystallization from MeOH/Acetone/H₂O (14 mL/46mL/0.325 mL) gave 2.1 g = 35% of analytically pure S(+)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol fumarate hydrate salt. Rotation: -6.56° (in methanol) for the fumarate hydrate. +9.07° (in

methanol) for the free base. Anal. Calcd: C, 60.43; H, 7.86; N, 3.52. Found: C, 60.47; H, 7.51; N, 3.32.

Example No. 6

5 **S(+)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol fumarate**
hydrate salt

Under gentle N₂ stream 60% NaH (35 mmols = 1.4 g) was washed with hexane, collected by filtration and transferred into a 250 mL 3 neck flask. DMF (20mL) was added into the flask to cover the sodium hydride and the suspension cooled to 10°C. Under stirring a solution of ethanthiol (32.40 mmols = 2.075 g = 2.47 mL) in DMF (5 mL) was added dropwise over 40 min. During the addition of the ethanthiol solution, foaming was noted in flask. Stirring was continued at 20°C for 1 1/2 h and the starting material S(+)-1-[2-(dimethylamino)-1-(4-methoxyphenyl)-ethyl]-cyclohexanol (12.5 mmols = 3.46 g) was added as a solid over 5 min. The reaction mixture was heated up slowly to 150°C over 30 min and stirred at this temperature for another hour. After this time the yellow brownish colored reaction mixture was rapidly cooled to 25°C and quenched by adding it into a flask containing H₂O (90 mL). The mixture was charcoaled and filtered through celite. The cake was washed with H₂O (1 × 25mL) and 1N NaOH (1 × 50mL). At this point the pH of the clear yellow colored solution was 13. This was extracted with toluene (1 × 60 mL), followed by hexane (1 × 60mL). Under stirring at room temperature it was neutralized to pH = 9 with conc. HCl (3 mL). The resulting suspension was cooled at 5°C, stirred for 1 h and the white solid was collected by filtration, dried in air overnight to give 2.4 g = 72% of the phenol (mp 230-232°C). This compound (8.35 mmols = 2.2 g) and fumaric acid (10.023 mmols = 1.163 g) were dissolved in hot methanol/acetone (1:3) mixture (48 mL) and filtered leaving a clear light yellow colored solution. Under stirring at room temperature H₂O (0.2 mL) was added to the solution. The solution became cloudy immediately. Stirring was resumed for 2 h and the resulting white solid was collected by filtration, washed with acetone (2 × 50 mL) and dried at 35°C in a vacuum oven overnight to give 2.4 g of analytically pure compound.

- 12 -

Tests were conducted to examine the effects of these compounds at 5-HT₂ receptor sites and on monamine uptake.

5

METHODS

Male Sprague-Dawley rats (180-260 g, Charles River) were used in all neurochemical assays. Rats were housed in temperature-controlled quarters on a 12 hr light/12 hr dark cycle with free access to food and water.

10

Neurotransmitter uptake inhibition

Uptake experiments were performed using a crude synaptosomal preparation made from the brain tissue of adult male Sprague-Dawley rats. The cortex of 1 rat for
15 NE and 5-HT uptake was removed on ice and homogenized in 20 volumes of 0.32 M sucrose/g tissue weight using a Potter-Elvehjem teflon homogenizer (3 strokes at 840 rpm). The homogenate was then centrifuged for 12 minutes at 1,000 × g at 0-4°C. The resulting supernatant was decanted into a chilled glass beaker and kept on ice until assayed. Protein concentration was determined by the method of Lowry et al.
20 (1).

For these experiments, all compounds were run in duplicate in concentrations of 0.003-30.0 μM. Each tube received buffer (790 μl in drug tubes, 800 μl in control tubes), 10 μl of drug or standard (0.1 μM DMI for NE uptake and 3.0 μM zimelidine
25 HCl for 5-HT uptake), 100 μl isotope (0.1 μM ³H-NE and 0.05 μM ¹⁴C-5-HT), and 100 μl tissue. Tubes were incubated at 37°C for 6 minutes. Incubation was terminated by the addition of 2.5 ml buffer followed by vacuum filtration using a Brandel filtration manifold with Whatman GF/B glass fiber filters and a second wash with 2.5 ml buffer. Filters were added to 10 ml Hydrofluor, shaken for 15 minutes
30 and counted in a Packard 460CD scintillation counter equipped with dual-label dpm data reduction. Results were expressed as pmol uptake/mg protein/min. IC₅₀'s for

- 13 -

uptake inhibition were calculated by linear regression of logit [% of active uptake] vs. log [concentration of test drug].

Results

5

O-Desmethyl venlafaxine, 4-[1-(2-dimethylamino)-2-(1-hydroxycyclohexyl)-ethyl]-phenol, and its S(+) and R(-) enantiomers were tested for their ability to inhibit NE and 5-HT neurotransmitter uptake. O-Desmethyl venlafaxine inhibited 5-HT uptake (IC_{50} = 0.20 μ M). Both enantiomers of O-Desmethyl venlafaxine were active in inhibiting 5-HT uptake with the (-) enantiomer being the more potent [(+)O-Desmethyl venlafaxine IC_{50} = 0.12 μ M; (-)O-Desmethyl venlafaxine = 0.06 μ M]. Venlafaxine and O-Desmethyl venlafaxine also inhibited NE uptake (IC_{50} = 0.72 μ M and 75% inhibition at 0.3 μ M, respectively). Both enantiomers of O-Desmethyl venlafaxine also inhibited NE uptake [(+)O-Desmethyl venlafaxine IC_{50} = 0.72 μ M; (-) O-Desmethyl venlafaxine IC_{50} = 0.27 μ M]. The (-) enantiomer of O-Desmethyl venlafaxine was more potent in inhibiting NE uptake.

Pharmaceutical compositions and formulations containing the enantiomers described herein can be produced in the same fashion and containing the same dosages as those described in the art for venlafaxine hydrochloride. The pharmaceutical formulations or compositions of this invention include those having as an active ingredient the R(-) enantiomer of O-Desmethyl venlafaxine substantially free of S(+) O-Desmethyl venlafaxine. This invention also includes formulations in which an active ingredient is the S(+) enantiomer of O-Desmethyl venlafaxine substantially free of R(-) O-Desmethyl venlafaxine. Each of these formulations also comprises one or more pharmaceutically useful excipients, carriers or adjuvants.

Formulations of the present invention may be produced using the S or R enantiomer of O-Desmethyl venlafaxine, or a pharmaceutically acceptable salt or salt hydrate thereof, in the same fashion as described for venlafaxine formulations in U.S. Patent Nos. 5,530,013 (Husbands et al.) and 5,506,270 (Upton et al.), both of which are incorporated herein by reference.

- 14 -

Preferred oral extended release formulations of this invention are comprised of the active enantiomer in admixture with microcrystalline cellulose and hydroxypropylmethylcellulose. Formed as beads or spheroids, the drug containing
5 formulation is coated with a mixture of ethyl cellulose and hydroxypropylmethyl cellulose to provide the desired level of coating, generally from about two to about twelve percent on a weight/weight basis of final product or more preferably from about five to about ten percent (w/w) , with best results obtained at from about 6 to about 8 percent (w/w). More specifically, the extended release spheroid formulations
10 of this invention comprise from about 30 to 40 percent R(-) O-desmethyl venlafaxine, from about 50 to about 70 percent microcrystalline cellulose, NF, from about 0.25 to about 1 percent hydroxypropylmethylcellulose, USP, and from about 5 to about 10 percent film coating, all on a weight/weight basis. And preferably, the spheroid formulations contain about 35 percent active ingredient, about 55 to 60 percent
15 microcrystalline cellulose NF (Avicel® PH101), about one half percent hydroxypropyl methylcellulose 2208 USP (K3, Dow, which has a viscosity of 3 cps for 2% aqueous solutions, a methoxy content of 19–24% and a hydroxypropoxy content of 4–13%), and from about 6 to 8 percent film coating.

20 The film coating is comprised of 80 to 90 percent of ethyl cellulose, NF and 10 to 20 percent hydroxypropyl methylcellulose (2910), USP on a weight/weight basis. Preferably the ethyl cellulose has a ethoxy content of 44.0–51% and a viscosity of 50 cps for a 5% aqueous solution and the hydroxypropylmethylcellulose is USP 2910 having a viscosity of 6 cps at 2% aqueous solution with a methoxy content of
25 28–30% and a hydroxypropoxy content of 7–12%. The ethyl cellulose used herein is Aqualon HG 2834.

Other equivalents of the hydroxypropylmethylcelluloses 2208 and 2910 USP and ethyl cellulose, NF, having the same chemical and physical characteristics as the
30 proprietary products named above may be substituted in the formulation without changing the inventive concept. Important characteristics of suitable hydroxypropyl-methylcelluloses include a low viscosity, preferably less than 10 cps and more preferably 2-5 cps, and a gel temperature above that of the temperature of the extrudate during extrusion. As explained below, these and other characteristics which
35 enable the extrudate to remain moist and soft (pliable) are preferred for the

- 15 -

hydroxypropylmethylcellulose. In the examples below, the extrudate temperature was generally 50-55°C.

Specific examples of extended release compositions of this invention include
5 the following.

Formulation Example 1.

A mixture of 44.8 parts (88.4 % free base) of O-desmethyl venlafaxine or a
10 salt or hydrate thereof, such as the fumarate hydrate salt, 74.6 parts of the microcrystalline cellulose, NF, and 0.60 parts of hydroxypropylmethyl cellulose 2208, USP, can be blended with the addition of 41.0 parts water. The plastic mass of material is then extruded, spheronized and dried to provide uncoated drug containing
15 spheroids.

Stir 38.25 parts of ethyl cellulose, NF, HG2834 and 6.75 parts of
hydroxypropyl methylcellulose 2910, USP in a 1:1 v/v mixture of methylene chloride
and anhydrous methanol until solution of the film coating material is complete.

20 To a fluidized bed of the uncoated spheroids apply 0.667 parts of coating solution per part of uncoated spheroids to obtain extended release, film coated spheroids having a coating level of 3%.

The spheroids can then be sieved to retain the coated spheroids of a particle
25 size between 0.85 mm to 1.76 mm diameter. These selected film coated spheroids are filled into hard gelatin capsules conventionally.

Formulation Example 2.

30 Same as for Example 1 except that 1.11 parts of the film coating solution per part of uncoated spheroids is applied to obtain a coating level of 5%.

Formulation Example 3.

35 Same as for Example 1 except that 1.33 parts of the film coating solution is applied to 1 part of uncoated spheroids to obtain a coating level of 6%.

- 16 -

Formulation Example 4.

5 Same as for Example 1 except that 1.55 parts of the film coating solution is applied to 1 part of uncoated spheroids to obtain a coating level of 7%.

10 One preferred extended release formulation of this invention comprises those of the active ingredient in spheroids comprised of microcrystalline cellulose and, optionally, hydroxypropylmethylcellulose coated with a mixture of ethyl cellulose and hydroxypropyl methyl cellulose. Preferably, the spheroids are comprised of about 30% to 40% O-desmethyl venlafaxine hydrochloride by weight, about 50% to about 70% microcrystalline cellulose, NF, by weight, and from about 0.25% to about 1% by weight of hydroxypropylmethylcellulose, USP, and coated with from about 2% to about 12% of total weight of film coating comprised of from about 80% to about 90% by weight of film coating of ethyl cellulose, NF, and from about 10% to about 20% by weight of film coating of hydroxypropylmethylcellulose, USP.

20 A specific extended release formulation according to the paragraph above is wherein the spheroids are composed of about 37% by weight of the O-desmethyl venlafaxine enantiomer, about 0.5% by weight of hydroxypropylmethylcellulose 2208, and about 62% by weight of microcrystalline cellulose. Another set of preferred compositions of this type are those wherein the film coating is comprised of ethyl cellulose (4.81% of total weight) and hydroxypropylmethylcellulose (0.85% of total weight). In another such composition the film coating comprises 6- 8% by weight of total weight, such as a film coating comprised of ethyl cellulose (2.48% of total weight) and hydroxypropylmethylcellulose (0.437% of total weight).

30 Yet another composition according to this invention are those wherein the film coating composition is comprised of ethyl cellulose having a 44.0-51.0% content of ethoxy groups and hydroxypropylmethylcellulose having a methoxy content of 28.0-30.0% and a hydroxypropoxy group content of 7.0-12.0%. Film coating compositions of this type may be comprised of about 85% by total weight of film coating of ethyl cellulose having a 44.0-51.0% content of ethoxy groups, and about 15% by total weight of film coating of hydroxypropylmethylcellulose having a methoxy content of 28.0-30.0% and a hydroxypropoxy group content of 7.0-12.0%. A more specific film coating composition of this sort is comprised of 85% by weight

- 17 -

of ethyl cellulose type HG 2834 and 15% by weight of hydroxypropylmethylcellulose type 2910.

Another extended release formulation for once daily administration of this invention comprises the O-desmethyl venlafaxine enantiomer, or a salt or hydrate thereof, which comprises spheroids containing 37.3% O-desmethyl venlafaxine enantiomer, 62.17% microcrystalline cellulose and 0.5% hydroxypropylmethylcellulose type 2208, coated with a quantity of a mixture comprised of 85% ethyl cellulose type HG 2834 and 15% hydroxypropyl-methylcellulose type 2910 sufficient to give coated spheroids having a dissolution profile which gives the desired release rate over a 24 hour period.

A further extended release formulation of this invention is manufactured such that the spheroids are comprised of about 6% to 40% active compound by weight, about 50% to about 940% microcrystalline cellulose, NF, by weight, and, optionally, from about 0.25% to about 1% by weight of hydroxypropylmethylcellulose, USP, and coated with from about 2% to about 12% of total weight of film coating comprised of from about 80% to about 90% by weight of film coating of ethyl cellulose, NF, and from about 10% to about 20% by weight of film coating of hydroxypropylmethylcellulose, USP. A preferred subset of these extended release formulations are those wherein the spheroids are composed of about 8.25% by weight of active compound, or a pharmaceutically acceptable salt or hydrate thereof, and about 91.75% by weight of microcrystalline cellulose, with a coating of from 3 to 5 % by weight of the total weight. Another preferred subset or group are those formulations wherein the spheroids are composed of about 16.5% by weight of active drug agent and about 83.5% by weight of microcrystalline cellulose, with a coating of from 4 to 6 % by weight of the total weight.

In other pharmaceutical compositions and formulations of this invention, the active ingredient comprises venlafaxine hydrochloride combined with the O-desmethyl enantiomer, with the non-active ingredients being those described herein or in other formulations for venlafaxine hydrochloride known in the art.

Uses of these extended release formulations may be described as a method for providing a therapeutic blood plasma concentration of active drug compound(s) over a 24 hour period with diminished incidences of nausea and emesis which comprises

- 18 -

administering orally to a patient in need thereof, an encapsulated, extended release formulation that provides a peak blood plasma level of active agent in from about four to about eight hours, said formulation containing O-Desmethyl venlafaxine, or a salt or salt hydrate thereof, as the active ingredient. The methods are also useful for

5 eliminating the troughs and peaks of drug concentration in a patients blood plasma attending the therapeutic metabolism of plural daily doses of active ingredient(s) which comprises administering orally to a patient in need thereof, an encapsulated, extended release formulation that provides a peak blood plasma level of the active agent in from about four to about eight hours, said formulation containing O-

10 Desmethyl venlafaxine, or a salt or salt hydrate thereof, as the active ingredient.

What is Claimed:

1. R(-)-4-[2-(Dimethylamino-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof, substantially free of S(+)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof.

2. S(+)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof, substantially free of R(-)-4-[2-(Dimethylamino-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof.

3. A pharmaceutical composition comprising one or more pharmaceutically acceptable carriers and a pharmaceutically effective amount of R(-)-4-[2-(Dimethylamino-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof, substantially free of S(+)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof.

4. A pharmaceutical composition comprising one or more pharmaceutically acceptable carriers and a pharmaceutically effective amount of S(+)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof, substantially free of R(-)-4-[2-(Dimethylamino-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof.

5. A method of treatment of depression in a mammal, the method comprising administering to a mammal in need thereof a pharmaceutically effective amount of R(-)-4-[2-(Dimethylamino-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof, substantially free of S(+)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof.

- 20 -

6. A method of treatment of depression in a mammal, the method comprising administering to a mammal in need thereof a pharmaceutically effective amount of S(+)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof, substantially free of R(-)-4-[2-(Dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol, or a pharmaceutically acceptable salt or salt hydrate thereof.

7. A process for the preparation of (R)-(-)-4-[2-(dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol or a pharmaceutically acceptable salt or salt hydrate thereof, which comprises

- (a) separating the R-enantiomer from the S-enantiomer by a standard resolution technique; or
- (b) subjecting (R)-4-[2-(dimethylamino)-1-(1- R₁O cyclohexyl)ethyl]phenol or a salt thereof, wherein R₁ is a removable protecting group, to a reaction to remove the protecting group; and, if desired, converting (R)-(-)-4-[2-(dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol into a salt thereof or converting a salt of (R)-(-)-4-[2-(dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol into (R)-(-)-4-[2-dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol.

8. A process for the preparation of (S)-(+)-4-[2-(dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol or a pharmaceutically acceptable salt or salt hydrate thereof, which comprises

- (c) separating the S-enantiomer from the R-enantiomer by a standard resolution technique; or
- (d) subjecting (S)-4-[2-(dimethylamino)-1-(1- R₁O cyclohexyl)ethyl]phenol or a salt thereof, wherein R₁ is a removable protecting group, to a reaction to remove the protecting group; and, if desired, converting (S)-(+)-4-[2-(dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol into a salt thereof or converting a salt of (S)-(+)-4-[2-(dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol into (S)-(+)-4-[2-dimethylamino)-1-(1-hydroxycyclohexyl)ethyl]phenol.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/16388

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C215/52 A61K31/135 A61P25/24

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 00 32556 A (SEPRACOR INC) 8 June 2000 (2000-06-08) page 1, line 20 - line 28 page 4, line 14,17,26,29 page 4, line 34 - line 36 page 5, line 28 - line 33 page 7, line 1 - line 8 page 14, line 8 - line 22 page 20, line 8 - line 18 page 24, line 1 -page 25, line 26 claims ---	1-8
P,X	WO 00 32555 A (SEPRACOR INC) 8 June 2000 (2000-06-08) page 3, line 1 - line 32 claims --- -/--	1-8



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

13 September 2000

Date of mailing of the international search report

21/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

O'Sullivan, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/16388

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>YARDLEY J P ET AL: "2-PHENYL-2-(1-HYDROXYCYCLOALKYL)ETHYLAMIN E DERIVATIVES: SYNTHESIS AND ANTIDEPRESSANT ACTIVITY" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 33, 1990, pages 2899-2905, XP000891765 ISSN: 0022-2623 cited in the application examples 6,8; table 1 page 2903, synthesis of (6)</p>	1-6
A	<p>US 5 530 013 A (HUSBANDS G E MORRIS ET AL) 25 June 1996 (1996-06-25) cited in the application column 1, line 16 - line 54 column 3, line 4 - line 7</p>	1-4
A	<p>US 5 788 986 A (DODMAN NICHOLAS H) 4 August 1998 (1998-08-04) cited in the application page 19, table 1, venlafaxine claim 1</p>	1-4

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/16388

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 0032556	A	08-06-2000	NONE		
WO 0032555	A	08-06-2000	NONE		
US 5530013	A	25-06-1996	CA	2141774 A	15-08-1995
			EP	0667150 A	16-08-1995
			JP	7252143 A	03-10-1995
			SG	49680 A	15-06-1998
US 5788986	A	04-08-1998	US	5554383 A	10-09-1996
			WO	9631172 A	10-10-1996
			US	5762960 A	09-06-1998